arXiv:1207.6554v1 [quant-ph] 27 Jul 2012

Collision-model-based approach to non-Markovian quantum dynamics

F. Ciccarello[1], M. G. Palma[2], and V. Giovannetti[1]
[1]*NEST, Scuola Normale Superiore and Istituto Nanoscienze-CNR, Piazza dei Cavalieri 7, I-56126 Pisa, Italy*
[2]*NEST, Istituto Nanoscienze-CNR and Dipartimento di Fisica,*
Università degli Studi di Palermo, via Archirafi 36, I-90123 Palermo, Italy

(Dated: July 30, 2012)

We present a theoretical framework to tackle quantum non-Markovian dynamics based on a microscopic collision model (CM), where the bath consists of a large collection of initially uncorrelated ancillas. Unlike standard memoryless CMs, we endow the bath with memory by introducing inter-ancillary collisions between next system-ancilla interactions. Our model interpolates between a fully Markovian dynamics and the continuous interaction of the system with a single ancilla, i.e., a strongly non-Markovian process. We show that in the continuos limit one can derive a general master equation, which while keeping such features is guaranteed to describe an unconditionally completely positive and trace-preserving dynamics. We also show that, for a specific choice of the involved couplings, known behaviors of an atomic system decaying in a damped cavity arise naturally in a suitable regime.

PACS numbers: 03.65.Yz, 03.67.-a, 42.50.Lc

Introduction. In open system dynamics problems the focus is on a system "S" in contact with an external bath or environment. Typically, the goal is to seek a master equation (ME) where the degrees of freedom (DOFs) of S are the only explicit variables. Hence, the environmental interactions should be accounted for through an effective but *reliable* description. When it comes to quantum objects, this problem turns out to be especially thorny [1–4]. Within this context, "reliable" means that the ME to be worked out should give rise to a completely positive and trace-preserving (CPT) dynamics. It is well-assessed that Markovian, i.e., memoryless, environments are described by MEs in the so called Lindlab form [1] entailing unconditionally CPT dynamics. Markovianity is in most cases only an approximation, though: in general, the environment is not forgetful and there is indeed a broad variety of actual phenomena featuring strong *non-Markovian* (NM) effects [5]. Yet, a general systematic framework for describing these has not been developed to date, which is a topical issue of concern to a manifold of variegated research areas. Rather, many different approaches have been proposed [6]. Typically, they are to some extent regularly underpinned by phenomenological assumptions and/or approximations (testifying the formidable hurdles to cope with in such problems). As a consequence, non-CPT – namely unphysical – dynamics can turn out in certain parameter ranges [7]. Among these descriptive tools are the so called memory-kernel MEs, e.g. those in Refs. [8, 9]. These are integro-differential MEs featuring a history integral, where past states of S are weighted through a certain memory-kernel function. There exist regimes in which such MEs can fail to be CPT [10–16]. What is more, it was recently tested [17] whether MEs in Refs. [8, 9] are non-Markovian according to a non-Markovianity indicator proposed by Breuer *et al.* [18]. It turned out that this is null [17], which suggests that such MEs should rather be regarded as time-dependent Markovian. This means that when they entail a CPT dynamics this is anyway very close to the purely Markovian regime.

An interesting picture to describe the interaction of S with a reservoir is to use a microscopic *collision model* (CM) [19–24]. The bath is modeled as a large collection of non-interacting identical ancillas (each can be thought as a low-dimensional system even though this is not necessary). By hypothesis, S "collides" with each of these one at a time and, importantly, is not allowed to interact more than once with a given ancilla. Demonstrably, such a process gives rise to an irreversible dynamics for S corresponding to a Lindlab-type [1], i.e., Markovian, ME [22]. This can be expected since, as stressed, at each step S comes into contact with a fresh ancilla which is still in its initial state. Hence, there is no way for the bath to keep track of the system's past history. Although they are somewhat fictitious, latest research is unveiling the potential of CMs as effective theoretical tools for tackling open system dynamics [23, 24]. First, they are conceptually intuitive, hence potentially easier to cope with: a complex coupling to a large environment is decomposed as a succession of elementary interactions with its subparts. A key feature is that CMs lead to Lindlab-type MEs without demanding any approximation: in fact, only the passage to the continuous limit is needed [22]. This is in contrast to standard microscopic system-reservoir models [1], where Markovianity must be somehow forced through drastic assumptions such as the requirement of small coupling and short enough bath correlation time (Born-Markov approximation). Should such a feature be maintained in a NM generalization of a CM, this would be quite appealing: as stressed above, approximations and phenomenological assumptions can lead to unphysical predictions. First progress along this line has been made very recently [23, 24]. A ME governing processes where many S systems interact with a bath in succession was worked out [23]. Rybar *et al.* [24] introduced a CM able to simulate any indivisible channel [1] (thus highly NM) when S is a single qubit [25], i.e., a two-level system. Memory was introduced by taking the bath ancillas initially in a nontrivial quantum-correlated state, whose form depends on the specific simulated channel.

Model. Physically, one would intuitively associate the bath ability to "remember" with some *dynamical* parameter some-



FIG. 1: (Color online) Sketch of the collisional model in the memoryless case (a) and in the NM one (b). Only the first steps are shown. The next ones are obtained through simple iteration.

how entering its interaction with S. In the spirit of CMs, a natural memory mechanism to devise is adding inter-ancillary (AA) collisions between next system-ancilla (SA) ones. This way, quantum information received from S can be conveyed across the bath and returned to S in next SA collisions. Here, we introduce a NM CM precisely built upon this idea. In the beginning (see Fig. 1) S collides with ancilla 1. In standard (Markovian) CMs, S-2 collision would then follow, then S-3 and so on [see Fig. 1(a)]. This way, each ancilla would still be in the initial state before colliding with S, thus fully "unaware" of previous collisions. In contrast, as sketched in Fig. 1(b), we assume that an extra AA collision between 1 and 2 occurs after S-1 but before S-2. Thereby, prior to its interaction with S, ancilla 2 will now be in a perturbed state in which information over past history of S is imprinted. The process proceeds by mere iteration: once S-2 collision is over, a 2-3 interaction follows, then S-3, 3-4 etc. Each collision, either SA or AA, is described by a CPT quantum map affecting the DOFs of the two involved particles. Specifically, the map for an SA collision involving the ith ancilla is defined as $\mathcal{U}_{Si}\sigma=\hat{U}_{Si}\sigma\hat{U}_{Si}^{\dagger}$, with $\hat{U}_{Si}=e^{-i\hat{H}_{Si}\tau}$ being a unitary operator which depends upon the collision time τ and the interaction Hamiltonian $\hat{H}_{Si}$ (we set $\hbar=1$ throughout). Instead the AA collision involving the ancilla i to the jth one is defined in terms of a CPT map $\mathcal{S}_{ij}$ which is modeled in one of two possible forms

$$\mathcal{S}_{ij}^{(c)}\sigma \;=\; \hat{V}_{ij}\sigma\hat{V}_{ij}^{\dagger} \quad \text{with} \quad \hat{V}_{ij}=\cos(g\tau)\mathbb{1}+i\sin(g\tau)\hat{S}_{ij}, \tag{1}$$

$$\mathcal{S}_{ij}^{(i)}\sigma \;=\; \cos(g\tau)^2\sigma+\sin(g\tau)^2\hat{S}_{ij}\sigma\hat{S}_{ij} \tag{2}$$

where g is a coupling rate while $\hat{S}_{ij}$ is the well-known swap operator [25] exchanging the states of i and j as $\hat{S}_{ij}\,|\varphi\rangle_i|\psi\rangle_j=|\psi\rangle_i|\varphi\rangle_j$ (tensor product symbols are omitted throughout). The transformation Eq. (1) is a unitary partial swap transferring information from i to j in terms of a two-body quantum gate. Map Eq. (2) is a sort of incoherent (non-unitary) analogue of Eq. (1) in that it accomplishes $i \to j$ information transfer in a probabilistic fashion: we will henceforth refer to maps (1) and (2) as coherent and incoherent partial swap, respectively, as indicated by the subscripts. We stress that in either case the map is CPT anyway. Let then ρ_0 be the initial of S, while each ancilla is initially in $|0\rangle\langle 0|$, where $|0\rangle$ is a given single-ancilla pure state (to fix the ideas, this can be thought as the ground state even though this is not necessary). The steps of our discrete process are defined in a such way that the nth step terminates when both (n-1)–n and S–n collisions are over (but step $n=1$ ending after S–1 collision). Hence, at the nth step the overall state is given by

$$\sigma_n=(\mathcal{U}_{Sn}\circ\mathcal{S}_{n,n-1}\circ\ldots\circ\mathcal{U}_{S2}\circ\mathcal{S}_{2,1}\circ\mathcal{U}_{S1})\,\sigma_0 \tag{3}$$

where $\sigma_0=\rho_0\,|\mathbf{0}\rangle_B\langle\mathbf{0}|$ is the system-bath initial state [26], where $|\mathbf{0}\rangle_B=|0\rangle_1|0\rangle_2\cdots$, while "$\circ$" represents the super-operator composition and will be henceforth omitted. In what follows, we shall be concerned with two cases, dubbed "(a)" and "(b)", respectively. In (a), we take the super-operators $\mathcal{S}_{ij}=\mathcal{S}_{ij}^{(i)}$ of Eq. (3) as given by Eq. (2) but leave $\mathcal{U}_{Si}$, i.e., $\hat{U}_{Si}$, unspecified. In (b), instead, we set $\mathcal{S}_{ij}=\mathcal{S}_{ij}^{(c)}$ [see Eq. (1)] but assume that $\hat{U}_{Si}$ corresponds to a Jaynes-Cummings-like S–i coupling [27].

(a): Incoherent partial swap. Map $\mathcal{S}_{ij}$ is given by Eq. (2). Let $\rho_n=\mathrm{Tr}_B(\sigma_n)$ be the nth -step state of S (the partial trace is over all the ancillas). We first show that ρ_n can be expanded in terms of all previous-step states of S *only*. By definition, $\hat{S}_{i,i+1}$ swaps the states of i and $(i+1)$, hence it transforms each $\hat{U}_{Si}$ as $\hat{S}_{i,i+1}\hat{U}_{Si}\,\hat{S}_{i,i+1}=\hat{U}_{S,i+1}$ (we recall that $\hat{S}_{ij}^{\dagger}\equiv\hat{S}_{ij}$). Equivalently, $\hat{S}_{i,i+1}\hat{U}_{Si}=\hat{U}_{S,i+1}\,\hat{S}_{i,i+1}$: a swap operator on the left of a $\hat{U}$-type one can jump to the right side provided that the ancillary index of $\hat{U}$ is increased by one unity. By construction [see Eq. (3)], $\sigma_2=\mathcal{U}_{S2}\mathcal{S}_{12}^{(i)}\sigma_1$, i.e., $\sigma_2=c^2\hat{U}_{S2}\,\sigma_1\hat{U}_{S2}^{\dagger}+s^2\hat{U}_{S2}S_{12}(\hat{U}_{S1}\,\sigma_0\hat{U}_{S1}^{\dagger})S_{12}\hat{U}_{S2}^{\dagger}$, where $c=\cos(g\tau)$, $s=\sin(g\tau)$ [cf. Eq. (2)] and we have replaced (only in the latter term) $\sigma_1=\hat{U}_{S1}\,\sigma_0\hat{U}_{S1}^{\dagger}$ [see Fig. 1(b)]. Operator $\hat{U}_{S1}$ can be eliminated as follows. We use $\hat{S}_{12}\hat{U}_{S1}=\hat{U}_{S,2}\,\hat{S}_{12}$ alongside $\hat{S}_{12}\sigma_0=\rho_0\hat{S}_{12}|\mathbf{0}\rangle_B\langle\mathbf{0}|\equiv\sigma_0$. This yields $\sigma_2=c^2\hat{U}_{S2}\,\sigma_1\hat{U}_{S2}^{\dagger}+s^2\hat{U}_{S2}^2\,\sigma_0\left(\hat{U}_{S2}^{\dagger}\right)^2$, where only $\hat{U}_{S2}$ now appears and all the states for $n<2$ are featured. For an arbitrary $n\geq 2$, this is generalized by induction [28] as

$$\sigma_n=c^2\sum_{j=1}^{n-1}s^{2(j-1)}\hat{U}_{Sn}^{j}\,\sigma_{n-j}\left(\hat{U}_{Sn}^{\dagger}\right)^j+s^{2(n-1)}\hat{U}_{Sn}^{n}\,\sigma_0\left(\hat{U}_{Sn}^{\dagger}\right)^n \tag{4}$$

where the coefficients fulfill the normalization constraint $c^2\sum_{j=1}^{n-1}s^{2(j-1)}+s^{2(n-1)}=1$. The form of Eq. (4) is attractive since, as we show next, a similar expansion for $\rho_n=\mathrm{Tr}_B\sigma_n$ holds. Each term, indeed, features a power of $\hat{U}_{Sn}$, involving the nth ancilla *only*, and a σ_{n-j}. Hence, upon trace over B the latter becomes the corresponding ρ_{n-j}

$$\rho_n=c^2\sum_{j=1}^{n-1}s^{2(j-1)}\mathcal{E}_j\,\rho_{n-j}+s^{2(n-1)}\mathcal{E}_n\,\rho_0\,. \tag{5}$$

In Eq. (5), $\mathcal{E}_j$ ($j=0,1,...$) are CPT maps on S defined as

$$\mathcal{E}_j\,\rho=\sum_{\nu}{}_n\langle\nu|\,\hat{U}_{Sn}^{j}\,|0\rangle_n\,\rho\,\left({}_n\langle\nu|\,\hat{U}_{Sn}^{j}\,|0\rangle_n\right)^{\dagger} \tag{6}$$

where each Kraus operator ${}_n\langle\nu|\,\hat{U}^{j}_{S\,n}\,|0\rangle_n$ is independent of n since so is the form of $\hat{U}_{S\,n}$ and the ancillas are all identical ($\{|\nu\rangle\}$ is a basis for the single-ancilla Hilbert space). Interestingly, the structure of Eq. (5) shares features with the discrete model used by Shabani and Lidar [9] to derive their NM ME. Two major differences occur, though. First, Eq. (5) cannot be written as a single sum due to the missing c^2 factor in the last term, which in fact means that here we deal with a time-inhomogeneous memory-kernel function (MKF). Second, map $\mathcal{E}_j$ is in general strongly NM since, due to $\hat{U}^{j}_{S\,i}=e^{-i\hat{H}_{Si}(j\tau)}$ (see *Model*), it effectively describes the continuous coherent interaction of S with a *single* ancilla. Indeed, our model interpolates between two extreme regimes depending on the value of s. When $s=0$, AA collisions are absent [cf. Eq. (2)]: Eq. (5) reduces to $\rho_n=\mathcal{E}_1\rho_{n-1}$ and we retrieve a standard Markovian CM [19–22]. Quite differently, for $s=1$, Eq. (5) yields $\rho_n=\mathcal{E}_n\rho_0$, i.e., S behaves as if it interacts with a single ancilla all the time. This can be seen by noting that for $s=1$ Eq. (2) reduces to a perfect swap: once S has undergone a τ-long interaction with i, the final state of i is fully transferred to $i+1$ (with i returning to $|0\rangle_i$). Our model thus interpolates between a fully Markovian and a strongly NM regime with s behaving as a sort of knob tuning the bath memory. Our next goal is to work out the ME corresponding to Eq. (5) in the continuous limit: We show that it likewise interpolates between the two opposite limits depicted above and is CPT in any case.

By subtracting from Eq. (5) the analogous identity for $n-1$, the variation of ρ_n between two next steps $\Delta\rho_n=\rho_n-\rho_{n-1}$ reads

$$\Delta\rho_n=c^2\textstyle\sum_{j=1}^{n-2}\,s^{2(j-1)}\mathcal{E}_j\Delta\rho_{n-j}+c^2s^{2(n-1)}\mathcal{E}_{n-1}\,\rho_1+\Delta\left(s^{2(n-1)}\mathcal{E}_n\right)\rho_0.\tag{7}$$

We introduce the elapsed time $t=n\tau$, which for $\tau\simeq 0$ and $n\gg 1$ becomes a continuous variable. Analogously, $t'=j\tau$. The sine powers in Eq. (7) can be expressed as a decaying exponential as $s^{2j}=(s^{2\frac{1}{\tau}})^{j\tau}=e^{-\Gamma t'}$, where $\Gamma=-2\ln s/\tau$ is an effective *memory rate*. Clearly, in the continuous limit τ must be the shortest time scale, hence $\Gamma\tau\ll 1$ and $c^2=1-s^2=1-e^{-\Gamma\tau}\simeq\Gamma\tau$. For vanishing τ, $\Delta\rho_n/\tau\to d\rho/dt$ and $\Delta\rho_{n-j}/\tau\to\partial\rho(t-t')/\partial(t-t')$. Thus once Eq. (7) is divided by τ, after a few straightforward technical steps [30] we end up with the ME

$$\frac{d\rho}{dt}=\Gamma\int_0^t dt' e^{-\Gamma t'}\mathcal{E}(t')\frac{\partial\rho(t-t')}{\partial(t-t')}+e^{-\Gamma t}\frac{d\mathcal{E}(t)}{dt}\rho_0 \tag{8}$$

where $\mathcal{E}(t)$ is the continuous analogue of Eq. (6). This is an integro-differential equation in $\rho(t)$ featuring a history integral term with an associated MKF $\Gamma e^{-\Gamma t'}$ and, notably, a term $\sim\rho_0$. The latter, which stems from the formerly discussed inhomogeneity of the discrete MKF in Eq. (5), is a strong signature of NM behavior. Indeed, for $\Gamma\to 0$ it is the only term surviving in Eq. (8) since $\partial_t\rho\to\partial_t\mathcal{E}\rho_0$, i.e., $\rho(t)\to\mathcal{E}(t)\rho_0$. Next, we derive the solution of Eq. (8) $\rho(t)=\Lambda(t)\rho_0$ and prove that the dynamical map [1] $\Lambda(t)$ is *always* CPT [$\Lambda(0)=\mathcal{I}$ with $\mathcal{I}$ the identity superoperator]. Evidently, $\Lambda(t)$ obeys Eq. (8) under the formal replacement $\rho\to\Lambda$. By taking the Laplace transform (LT) of such equation, this is easily solved as [30]

$$\tilde{\Lambda}(s)=\frac{\tilde{\mathcal{E}}(s+\Gamma)}{\mathcal{I}-\Gamma\,\tilde{\mathcal{E}}(s+\Gamma)}\tag{9}$$

where $\tilde{\Lambda}(s)$ and $\tilde{\mathcal{E}}(s)$ are the LTs of $\Lambda(t)$ and $\mathcal{E}(t)$, respectively [Eq. (9) is well-defined since the numerator and denominator commute]. Expanding Eq. (9) in powers of Γ gives $\tilde{\Lambda}(s)=\sum_{k=1}^{\infty}\left[\tilde{\mathcal{E}}(s+\Gamma)\right]^k\Gamma^{k-1}$, whose inverse LT is

$$\Lambda(t)=\mathcal{L}^{-1}[\tilde{\Lambda}(s)](t)=\sum_{k=1}^{\infty}\Gamma^{k-1}\;\mathcal{L}^{-1}[\tilde{\mathcal{E}}^k(s+\Gamma)](t)\,.\tag{10}$$

Basic properties of LT allow to immediately calculate the inverse LT within braces as [30]

$$\begin{aligned}\mathcal{L}^{-1}[\tilde{\mathcal{E}}^k(s+\Gamma)] \;=\; & e^{-\Gamma t}\int_0^t dt_1\int_0^{t_1}dt_2\cdots\int_0^{t_{k-2}}dt_{k-1}\\ & \times\mathcal{E}(t_{k-1})\mathcal{E}(t_{k-2}-t_{k-1})\cdots\mathcal{E}(t-t_1).\end{aligned}\tag{11}$$

We have thus expressed $\Lambda(t)$ as a weighted series of multiple auto-convolutions of the CPT map $\mathcal{E}(t)$. Being a composition of CPT maps, each convolution [the integrand in Eq. (11)] is CPT itself. Also, it is multiplied by a positive coefficient [cf. Eqs. (10) and (11)], which yields complete positivity of map $\Lambda(t)$. Moreover, the state obtained by applying the *integrand* in Eq. (11) to ρ_0 has evidently unitary trace. As is easily checked [30], this entails $\mathrm{Tr}\,[\Lambda(t)\rho_0]=1$. We conclude that map $\Lambda(t)$ is *always* CPT. The last remaining task is to prove that, in line with Eq. (5) for $s=0$, the Markovian behavior arises from Eq. (8) for $\Gamma\to\infty$. Indeed, Eq. (8) is such that for Γ large enough we can approximate $\mathcal{E}(t)\simeq\mathcal{I}+\mathcal{F}t$, where $\mathcal{F}=\dot{\mathcal{E}}(0)$. Under LT, this becomes $\tilde{\mathcal{E}}(s)=1/(s+\Gamma)+\mathcal{F}/(s+\Gamma)^2$, which once plugged into Eq. (9) and in the limit of $\Gamma\to\infty$ yields $\tilde{\Lambda}(s)=(s+\Gamma+\mathcal{F})/[s^2+\Gamma(s-\mathcal{F})]\big|_{\Gamma\to\infty}=1/(s-\mathcal{F})$. By transforming back, we end up with $\Lambda(t)=e^{\mathcal{F}t}$ entailing that the semigroup property is fulfilled and thus, necessarily, $\mathcal{F}$ is a Lindlabian superoperator [1] with Eq. (8) reducing to the Lindlab-form $\dot{\rho}=\mathcal{F}\rho$.

To conclude this section, we point out that if one sets $\mathcal{E}(t)=e^{\mathcal{F}t}$, where $\mathcal{F}$ is a Lindlabian, then it is immediate to check that $\Lambda(t)=e^{\mathcal{F}t}$ is the *exact* solution of Eq. (8) for *any* Γ. This further marks the difference with the Shabani-Lidar ME [9], where instead the fact that $\Lambda(t)=e^{\mathcal{F}t}$ is not a solution in that context is exploited in a perturbative way to deliver the conclusions. This further strengthens the importance of the last term in Eq. (8). Also note that our ME agrees with the general form predicted by Nakajima and Zwanzig [1] provided that the corresponding $t-t'$-dependent memory-kernel superoperator exhibits a discontinuity at $t=t'$.

(b): Coherent partial swap. Each AA collision is now described by the *unitary* map in Eq. (1). Let $\hat{\mathrm{U}}_n$ be the (discrete) overall evolution unitary operator such that $\sigma_n=\hat{\mathrm{U}}_n\sigma_0\hat{\mathrm{U}}_n^{\dagger}$ and $\hat{\mathrm{U}}_0=\mathbb{1}$. From Eqs. (1) and (3), $\hat{\mathrm{U}}_1=\hat{U}_{S\,1}$ while $\hat{\mathrm{U}}_{n\geq 2}=\hat{U}_{S\,n}\hat{V}_{n,n-1}\cdots\hat{U}_{S\,2}\hat{V}_{21}\hat{U}_{S\,1}$. By using the properties of the swap $\hat{S}_{ij}$ alongside Eq. (1) and $\sigma_0=\rho_0\,|\mathbf{0}\rangle_B\langle\mathbf{0}|$, a reasoning basically

analogous to the one leading to Eq. (4) – with due replacements – allows to express $\hat{\mathrm{U}}_{n\geq 2}$ *effectively* as [29]

$$\hat{\mathrm{U}}_n = c\sum_{j=1}^{n-1}(is)^{j-1}\hat{U}_{Sn}^{j}\hat{\mathrm{U}}_{n-j} + (is)^{n-1}\hat{U}_{Sn}^{n}\hat{\mathrm{U}}_0 \qquad (12)$$

which upon use of $\sigma_n = \hat{\mathrm{U}}_n\sigma_0\hat{\mathrm{U}}_n^\dagger$ yields $\sigma_n = c^2\sum_{j=1}^{n-1} s^{2(j-1)}\hat{U}_{Sn}^{j}\,\sigma_{n-j}\hat{U}_{Sn}^{j\dagger} + s^{2(n-1)}\hat{U}_{Sn}^{n}\,\sigma_0\hat{U}_{Sn}^{n\dagger}$ + o.d.t. where "o.d.t." stands for off-diagonal terms. Note that the diagonal part coincides with Eq. (4), hence the effect of replacing map $\mathcal{S}_{ij}^{(i)}$ with $\mathcal{S}_{ij}^{(c)}$ is to introduce such o.d.t. Clearly, the trace over the bath now yields $\rho_n = c^2\sum_{j=1}^{n-1} s^{2(j-1)}\mathcal{E}_j\,\rho_{n-j} + s^{2(n-1)}\mathcal{E}_n\,\rho_0 +$ Tr_B(o.d.t.). Likewise, this differs from Eq. (5) due to the trace over off-diagonal terms. Yet, these are absent for $s=0$ and $s=1$ since in either case Eq. (12) reduces to a single term. Hence, our comments on these two regimes presented in the case (a) hold unchanged. A general treatment accounting for o.d.t. is demanding. However, the continuous-limit dynamics can be worked out exactly in the case that S is a qubit and the SA coupling has the form of an *XY isotropic interaction*, i.e., a situation routinely encountered in quantum optics problems. As in such a case the total number of excitations is conserved, each ancilla behaves as an effective qubit as well. Let $\{|0\rangle_{S(i)},|1\rangle_{S(i)}\}$ be a basis for $S(i)$. The SA Hamiltonian reads $\hat{H}_{Si} = \Omega\left(\hat{\sigma}_+\hat{S}_{i-} + \mathrm{H.c.}\right)$, where Ω is a coupling rate while $\hat{\sigma}_+ = \hat{\sigma}_-^\dagger = |1\rangle_S\langle 0|$ and $\hat{S}_{i+} = \hat{S}_{i-}^\dagger = |1\rangle_i\langle 0|$ are the usual spin-1/2 ladder operators. The most general initial state of S reads $\rho_0 = (1-p)\,|0\rangle_S\langle 0| + p\,|1\rangle_S\langle 1| + (r\,|0\rangle_S\langle 1| + \mathrm{H.c.})$, where $0\leq p\leq 1$ and $|r|^2\leq p(1-p)$. To calculate $\sigma_n = \hat{\mathrm{U}}_n\,\rho_0\,|\mathbf{0}\rangle\langle\mathbf{0}|\,\hat{\mathrm{U}}_n^\dagger$, we use $\hat{\mathrm{U}}_n\,|0\mathbf{0}\rangle_{SB} = (c+is)^{n-1}\,|0\mathbf{0}\rangle_{SB} = e^{ig\tau(n-1)}|0\mathbf{0}\rangle_{SB}$, which follows from $\hat{H}_{Si}|0\mathbf{0}\rangle = 0$ and Eq. (1), and $\hat{\mathrm{U}}_n\,|1\mathbf{0}\rangle_{SB} = \alpha_n|1\mathbf{0}\rangle_{SB} + \beta_n|0\Psi\rangle_{SB}$, where α_n and β_n are complex coefficients and $|\Psi\rangle_B = \sum_i c_i\,|i\rangle_B$ is some one-excitation state of B ($|i\rangle_B$ is the state such that each ancilla is in $|0\rangle$ but the ith ancilla, which is in $|1\rangle$). Thus $\rho_n = \mathrm{Tr}_B\sigma_n = 1 - p|z_n|^2\,|0\rangle_S\langle 0| + p|z_n|^2\,|1\rangle_S\langle 1| + \{r\,z_n\,|0\rangle_S\langle 1| + \mathrm{H.c.}\}$ with $z_n = e^{ig\tau(n-1)}\alpha_n^*$. This is the well-known form of an amplitude-damping channel [31]. The dynamical map is thus ensured to be CPT and is fully specified by z_n. This fulfills the identity

$$z_n = c\sum_{j=1}^{n-1}\left[e^{i\left(g\tau-\frac{\pi}{2}\right)}s\right]^{j-1}\cos(\Omega j\tau)\,z_{n-j} + \left[e^{i\left(g\tau-\frac{\pi}{2}\right)}s\right]^{n-1}\cos(\Omega n\tau), \qquad (13)$$

which can be shown in a very few steps [30].

To pass to the continuous limit, similarly to the case of the incoherent swap we define $t=n\tau$ by setting $\tau\simeq 0$ and $n\gg 1$. Analogously, in Eq. (13) $j\rightarrow t'$. We now consider the regime of nearly perfect swap $g\tau\simeq\pi/2$ [cf. Eq. (1)], which entails $c = \cos(g\tau)\simeq\pi/2 - g\tau = \bar{g}\tau$, where we have defined an effective rate $\bar{g}$ that evidently fulfills $\bar{g}\tau\ll 1$. Similarly to case (a), $s^{j-1}\simeq(s^{1/\tau})^{j\tau} = e^{-\gamma t'}$, where $\gamma\tau = -\ln s$. Using $s\simeq 1 - c^2/2$ we find $\gamma = \bar{g}^2\tau/2$. We thus end up with the integral equation

$$z(t) = i\,\bar{g}\int_0^t dt'\,e^{-i\bar{g}t'}e^{-\gamma t'}\cos(\Omega t')\,z(t-t') + e^{-i\bar{g}t}e^{-\gamma t}\cos(\Omega t) \qquad (14)$$

which is easily solved through the LT method [30] as

$$z(t) = \frac{e^{-\left[\gamma + i\left(\bar{g} - \sqrt{\Omega^2 + \bar{g}^2/4}\right)\right]t}}{2}\left[\frac{\bar{g}\left(-1 + e^{i2\sqrt{\Omega^2+\bar{g}^2/4}\,t}\right)}{2\sqrt{\Omega^2+\bar{g}^2/4}} + \left(1 + e^{i2\sqrt{\Omega^2+\bar{g}^2/4}\,t}\right)\right]. \qquad (15)$$

The squared absolute value of Eq. (15) is $|z(t)|^2 = \frac{e^{-2\gamma t}}{4\Omega^2+\bar{g}^2}\left[2\Omega^2 + \bar{g}^2 + 2\Omega^2\cos\left(2\sqrt{\Omega^2+\bar{g}^2/4}\,t\right)\right]$. Note that $0\leq|z(t)|^2\leq 1$ regardless of all the parameters showing that the dynamics remains CPT in the continuos limit. Also, $|z(t)|^2\equiv|\alpha(t)|^2$ is the excitation probability of S when it is initially in $|1\rangle_S$. For $\gamma = \bar{g} = 0$ (perfect swapping) we retrieve $|z(t)|^2 = \cos(\Omega t)^2$, as expected. Remarkably, in the regime $\Omega\gg\bar{g}$, $|z(t)|^2\simeq\exp[-2\gamma t]\cos(\Omega t)^2$ [indeed the integral term in Eq. (14) becomes negligible]. This coincides with the behavior of an atom decaying in a lossy single-mode cavity with damping rate γ predicted by a microscopically derived ME [32], which in turn is a more rigorous and accurate description than the standard phenomenological ME [33, 34]. Indeed, we have pointed out that for $s=1$ [cf. Eq. (1)] S behaves as if it is continuously interacting with a single ancilla. In the present case (b), the latter can be interpreted as the protected mode of a cavity on resonance with the atom, namely S. A Jaynes-Cummings dynamics [27] thus takes place. If the cavity is damped, leakage from the protected mode to the electromagnetic vacuum then occurs. In other words, from time to time the cavity mode can irreversibly lose the excitation received from the atom. In our CM, this corresponds to $s\neq 1$: from time to time, an AA collision fails to happen meaning that part of the excitation released to the bath will no longer be returned to S in next collisions.

Conclusions. We have introduced a NM microscopic CM, where the bath memory is added in a dynamical way through simple inclusion of inter-ancillary collisions. Each of these can be modeled as a CPT swapping operation. The model interpolates between two extreme situations: a fully Markovian regime (a standard memoryless CM is then retrieved) and a strongly NM one (corresponding to continuous interaction of the system with a single ancilla). We have shown the effectiveness of this framework in two different cases. For incoherent swapping, the resulting dynamics was shown to be described in the continuous limit by an exact ME holding for arbitrary system-ancilla coupling and regardless of the dimensionality of the involved particles. Such ME possesses two unusual – normally challenging – features. On the one hand, it encompasses both a fully Markovian regime and a strongly NM one as limiting cases. On the other hand, it is guaranteed to be unconditionally CPT. For coherent swapping, we have focused on the situation where S is a qubit coupled to each ancilla via a Jaynes-Cummings-like interaction. We have shown that in the continuous limit the corresponding dynamical map can be worked out exactly and is unconditionally CPT. Significantly, in a certain regime it retrieves known behaviors of an atomic system decaying in a damped cavity.

In conclusion, our findings witness the potential of a CM-based approach to tackle the daunting problem of a comprehensive and reliable description of NM dynamics.

We thank R. Fazio for comments and discussions and acknowledge support from the MIUR through the FIRB-IDEAS project RBID08B3FM.

Supplementary material

In this Supplementary Material, we supply some technical details related to the derivation of some properties discussed the paper's main text.

Derivation of Eq. (8)

When Eq. (7) is divided by τ and by using the limiting expressions discussed in the main text, the terms on the right-hand side in the continuous limit take the form

$$\frac{c^2\sum_{j=1}^{n-2} s^{2(j-1)}\mathcal{E}_j\left(\rho_{n-j}-\rho_{n-1-j}\right)}{\tau}\simeq\Gamma\int_0^t dt' e^{-\Gamma t'}\mathcal{E}(t')\frac{d\rho(t-t')}{d(t-t')}$$

$$\frac{\Delta(s^{2(n-1)}\mathcal{E}_n)}{\tau}\,\rho_0=\frac{(s^{2(n-1)}\mathcal{E}_n-s^{2(n-2)}\mathcal{E}_{n-1})}{\tau}\,\rho_0$$

$$\simeq\frac{e^{-\Gamma(t+2\tau)}\mathcal{E}_{t+\tau}-e^{-\Gamma(t+\tau)}\mathcal{E}(t)}{\tau}\,\rho_0=\frac{d}{dt}\Big(e^{-\Gamma t}\mathcal{E}(t)\Big)\rho_0$$

$$\frac{c^2 s^{2(n-1)}\mathcal{E}_{n-1}}{\tau}\,\rho_1\simeq\Gamma e^{-\Gamma t}\mathcal{E}(t)\,\rho_0$$

whereas the left-hand side of Eq. (7) clearly reduces to the time derivative of $\rho(t)$. Using these, Eq. (8) is immediately obtained.

Derivation of $\tilde{\Lambda}(s)$

By replacing $\rho(t) = \Lambda(t)\rho(0)$ in Eq. (8) and using that ρ_0 is arbitrary, the equation obeyed by $\Lambda(t)$ tuns out to be

$$\dot{\Lambda}(t) = \Gamma\int_0^t dt' \exp[-\Gamma t']\,\mathcal{E}(t')\dot{\Lambda}(t-t') + \exp[-\Gamma t]\,\dot{\mathcal{E}}(t) \qquad (16)$$

in addition to the requirement $\Lambda(0)=\mathcal{I}$. Upon Laplace transform (LT), the equation becomes

$$s\tilde{\Lambda}(s)-\mathcal{I}=\Gamma\tilde{\mathcal{E}}(s+\Gamma)[s\tilde{\Lambda}(s)-\mathcal{I}]+(s+\Gamma)\,\tilde{\mathcal{E}}(s+\Gamma)-\mathcal{I} \qquad (17)$$

where for s complex the LT is defined as

$$\tilde{F}(s)=\mathcal{L}\,[F(t)](s)=\int_0^\infty dt\; e^{-st}F(t) \qquad (18)$$

and we have used $\mathcal{E}(0)=\mathcal{I}$ (see main text).

By rearranging terms in Eq. (17)

$$[\mathcal{I}-\Gamma\,\tilde{\mathcal{E}}(s+\Gamma)][s\tilde{\Lambda}(s)-\mathcal{I}] = (s+\Gamma)\,\tilde{\mathcal{E}}(s+\Gamma)-\mathcal{I}\;, \qquad (19)$$

and hence

$$s\,[\mathcal{I}-\Gamma\,\tilde{\mathcal{E}}(s+\Gamma)]\,\tilde{\Lambda}(s) = s\,\tilde{\mathcal{E}}(s+\Gamma)\;. \qquad (20)$$

By simplifying s on both terms and introducing the inverse of $\mathcal{I}-\Gamma\,\tilde{\mathcal{E}}(s+\Gamma)$ we end up with Eq. (9).

Expansion of $\Lambda(t)$

The inverse LT of $\tilde{\mathcal{E}}(s+\Gamma)$ is $\mathcal{L}^{-1}[\tilde{\mathcal{E}}(s+\Gamma)] = e^{-\Gamma t}\mathcal{E}(t)$. Thereby, from a basic property of LT, the inverse transform of $\tilde{\mathcal{E}}^2(s+\Gamma)$ is the auto-convolution of $e^{-\Gamma t}\mathcal{E}(t)$, which reads

$$\begin{aligned}\mathcal{L}^{-1}[\tilde{\mathcal{E}}^2(s+\Gamma)] &= \int_0^t dt'\left[e^{-\Gamma t'}\mathcal{E}(t')\right]\left[e^{-\Gamma(t-t')}\mathcal{E}(t-t')\right]\\ &= e^{-\Gamma t}\int_0^t dt'\mathcal{E}(t')\mathcal{E}(t-t')\,. \qquad (21)\end{aligned}$$

The inverse LT of $\tilde{\mathcal{E}}^3(s+\Gamma)$ can be calculated as the convolution between Eq. (21) and $\mathcal{L}^{-1}[\tilde{\mathcal{E}}(s+\Gamma)]=\mathcal{E}(t)$, which yields

$$\mathcal{L}^{-1}[\tilde{\mathcal{E}}^3(s+\Gamma)]=e^{-\Gamma t}\int_0^t dt_1\int_0^{t_1} dt_2\;\mathcal{E}(t_2)\mathcal{E}(t_1-t_2)\mathcal{E}(t-t_1) \qquad (22)$$

Eq. (11), i.e., the case corresponding to $\tilde{\mathcal{E}}^k(s+\Gamma)$ for arbitrary k, then follows by mere induction.

Trace preservation of $\Lambda(t)$

As discussed in the main text, the integrand in Eq. (11) is a CPT map and thus, once applied to ρ_0, it yields a state having unitary trace (clearly, $\mathrm{Tr}\rho_0=1$). Hence,

$$\begin{aligned}\mathrm{Tr}\,[\Lambda(t)\rho_0] &= \sum_{k=1}^{\infty}\Gamma^{k-1}\,\mathrm{Tr}\left\{e^{-\Gamma t}\int_0^t dt_1\int_0^{t_1} dt_2\cdots\int_0^{t_{k-2}} dt_{k-1}\right\}\\ &= e^{-\Gamma t}\sum_{k=1}^{\infty}\Gamma^{k-1}\frac{t^{k-1}}{(k-1)!}=e^{-\Gamma t}e^{\Gamma t}=1\;. \qquad (23)\end{aligned}$$

Proof of Eq. (13)

By definition (see main text) z_n is related to α_n as $z_n = e^{ig\tau(n-1)}\alpha_n^*$. To derive α_n, we observe that owing to the assumed form of $\hat{H}_{S\,i}$ alongside Eq. (1) both SA and AA collisions conserve the total number of excitations. Thus, when $\rho_0 = |1\rangle_S\langle 1|$ the nth-step state is pure and necessarily reads $\hat{\mathrm{U}}_n\,|1\mathbf{0}\rangle_{S\,B} = \alpha_n|1\mathbf{0}\rangle_{S\,B}+\beta_n|0\Psi\rangle_{S\,B}$, where α_n and β_n are complex coefficients and $|\Psi\rangle_B=\sum_i c_i\,|i\rangle_B$ is a single-excitation bath state ($|i\rangle_B$ is the product bath state such that each ancilla is in $|0\rangle$ but the ith ancilla, which is in $|1\rangle$). Thus $\alpha_n=\langle 1\mathbf{0}|\,\hat{\mathrm{U}}_n\,|1\mathbf{0}\rangle$, which using Eq. (12) becomes $\alpha_n=\sum_{j=1}^n\mu_j\,\langle 1\mathbf{0}|\,\hat{U}_{S\,n}^{\,j}\hat{\mathrm{U}}_{n-j}\,|1\mathbf{0}\rangle$, where $\mu_{j\le n-1} = c(is)^{j-1}$, $\mu_n = (is)^{n-1}$. By diagonalizing $\hat{H}_{S\,n}$ it is immediately found that $\langle 1\mathbf{0}|_{S\,B}\hat{U}_{S\,n}^{\,j} = \cos(\Omega j\tau)\langle 1\mathbf{0}|_{S\,B} - i\sin(\Omega j\tau)\langle 0\,n|_{S\,B}$, where $\pm\Omega$ are the eigenvalues of $\hat{H}_{S\,n}$. As our process is such that the nth ancilla remains in $|0\rangle$ until the nth step, $|0\,n\rangle_{SB}$ does not overlap $\hat{\mathrm{U}}_{n-j}\,|1\mathbf{0}\rangle$ for $j\ge 1$. Therefore

$$\alpha_n=\sum_{j=1}^n\mu_j\,\langle 1\mathbf{0}|\,\hat{U}_{S\,n}^{\,j}\hat{\mathrm{U}}_{n-j}\,|1\mathbf{0}\rangle=\sum_{j=1}^n\mu_j\cos(\Omega j\tau)\,\alpha_{n-j} \qquad (24)$$

where $\alpha_0 = 1$. This immediately gives rise to Eq. (13) since $z_n=e^{ig\tau(n-1)}\alpha_n^*$.

Proof of Eq. (15)

Let $\tilde{z}(s)$ and $\tilde{f}(s)$ be the LTs of $z(t)$ and $f(t) = e^{-i\overline{g}t}e^{-\gamma t}\cos(\Omega t)$, respectively. Then the LT of Eq. (14) reads $\tilde{z}(s) = i\overline{g}\tilde{f}(s)\tilde{z}(s) + \tilde{f}(s)$, hence $\tilde{z}(s) = \tilde{f}(s)/[1 - i\overline{g}\tilde{f}(s)]$. The inverse LT of this is Eq. (15).

[1] H. P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford, Oxford University Press, 2002).

[2] U. Weiss, *Quantum Dissipative Systems*, 3rd ed. (World Scientific, Singapore, 2008).

[3] A. Rivas and S.F. Huelga, *Open Quantum Systems. An Introduction* (Springer, Heidelberg, 2011).

[4] H.-P. Breuer, arXiv: 1206.5346.

[5] See e.g.: J. Schliemann, A. Khaetskii, and D. Loss, J. Phys.: Condens. Matter **15**, R1809 (2003); M. Michel, G. Mahler, and J. Gemmer, Phys. Rev. Lett. **95**, 180602 (2005).

[6] Due to space constraints, our introduction is not intended to provide a comprehensive overview of the many different approaches proposed in the literature so far. We therefore focus on those more closely related to the present work.

[7] B. Vacchini and H. P. Breuer, Phys. Rev. A **81**, 042103 (2010).

[8] S. M. Barnett and S. Stenholm, Phys. Rev. A **64**, 033808 (2001).

[9] A. Shabani and D. A. Lidar, Phys. Rev. A **71**, 020101(2005).

[10] A. A. Budini, Phys. Rev. A **69**, 042107 (2004).

[11] S. Daffer *et al.*, Phys. Rev. A **70**, 010304(R) (2004).

[12] S. Maniscalco and F. Petruccione, Phys. Rev. A **73**, 12111 (2006); S. Maniscalco, Phys. Rev. A **75**, 062103 (2007).

[13] H. P. Breuer and B. Vacchini, Phys. Rev. E **79**, 041147 (2009). and

[14] J. Wilkie and Y. M. Wong, J. Phys. A **42**, 015006 (2009).

[15] S. Campbell et al., Phys. Rev. A **85**, 032120 (2012).

[16] D. Chruściński and A. Kossakowski, Phys. Rev. Lett. **97**, 20005 (2012); D. Chruściński and A. Kossakowski, Europhys. Lett. **97**, 20005 (2012).

[17] L. Mazzola, E.-M. Laine, H.-P. Breuer, S. Maniscalco, and J. Piilo, Phys. Rev. A **81**, 062120 (2010).

[18] H.-P. Breuer, E.-M. Laine, and J. Piilo, Phys. Rev. Lett. **103**, 210401 (2009).

[19] J. Rau, Phys. Rev. **129**, 1880 (1963).

[20] R. Alicki and K. Lendi, *Quantum Dynamical Semigroups and Applications*, Lecture Notes in Physics (Springer-Verlag, Berlin, 1987).

[21] M. Ziman *et al.*, Phys. Rev. A **65** , 042105 (2002); V. Scarani *et al.*, Phys. Rev. Lett. **88** , 97905 (2002)

[22] M. Ziman and V. Buzek, Phys. Rev. A **72**, 022110 (2005); M. Ziman, P. Stelmachovic, and V. Buzek, Open systems and information dynamics **12**, 81 (2005).

[23] V. Giovannetti and M. Palma, Phys. Rev. Lett. **108**, 040401 (2012); arXiv:1204.1836v1 [quant-ph], to appear in J. Phs. B.

[24] T. Rybar, S. N. Filippov, M. Ziman, and V. Buzek, arXiv:1202.6315.

[25] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, U. K., 2000).

[26] Our notation is such that states of S are denoted by ρ, while σ is used for states of the overall system (S and the bath).

[27] E. T. Jaynes and F. W. Cummings, Proc. IEEE **51**, 89 (1963).

[28] By construction, $\sigma_{n+1} = \mathcal{U}_{S,n+1}\mathcal{S}^{(i)}_{n,n+1}\sigma_n = c^2\hat{U}_{S,n+1}\,\sigma_n\hat{U}^{\dagger}_{S,n+1} + s^2\hat{U}_{S,n+1}S_{n,n+1}\sigma_n S_{n,n+1}\,\hat{U}^{\dagger}_{S,n+1}$. By replacing Eq. (4) in the second term and using $\hat{S}_{n,n+1}\hat{U}_{sn} = \hat{U}_{s,n+1}\hat{S}_{n,n+1}$ along with $\hat{S}_{n,n+1}\sigma_{n-j} = \sigma_{n-j}$ for $j \geq 1$, we end up with Eq. (4) for $n \rightarrow n+1$.

[29] In Eq. (12), we have kept the same notation for the evolution operators $\hat{\mathrm{U}}_n$. We point out, though, that Eq. (12) holds only for their *effective forms* given the initial state $\sigma_0 = \rho_0\,|\mathbf{0}\rangle_B\langle\mathbf{0}|$ (this is the only one considered in the present work).

[30] For technical details see supplementary material at [...].

[31] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, U. K.,2000).

[32] M. Scala *et al.*, Phys. Rev. A **75**, 013811 (2007).

[33] H.-J. Briegel and B.-G. Englert, Phys. Rev. A **47**, 3311 (1993).

[34] Significantly, at variance with the phenomenological ME [33], the ME in Ref. [32] treats as open system the bipartite one comprising both the atom and the cavity mode.